


One or a combination of symptoms can be warning signs of PH

The first warning sign may be a single kidney stone in children or recurrent stones in adults, and can include one or a combination of the following¹⁻¹¹:



Abbreviations: CKD, chronic kidney disease; ESRD, end-stage renal disease; UTIs, urinary tract infections.

1. Edvardsson VO, et al. *Pediatr Nephrol.* 2013;28(10):1923-1942. 2. Bhasin B, et al. *World J Nephrol.* 2015;4(2):235-244. 3. van Woerden CS, et al. *Nephrol Dial Transplant.* 2003;18(2):273-279. 4. Soliman NA, et al. *Nephrol Ther.* 2017;13(3):176-182. 5. Frishberg Y, et al. *Am J Nephrol.* 2005;25(3):269-275. 6. Johnson SA, et al. *Pediatr Nephrol.* 2002;17(8):597-601. 7. Garrelfs SF, et al. *Kidney Int.* 2019;96(6):1389-1399. 8. Belostotsky R, et al. *Am J Hum Genet.* 2010;87(3):392-399. 9. Harambat J, et al. *Kidney Int.* 2010;77(5):443-449. 10. Birtel J, et al. *Am J Ophthalmol.* 2019;206:184-191. 11. Mandrile G, et al. *Kidney Int.* 2014;86(6):1197-1204. 12. Estimated glomerular filtration rate (eGFR). National Kidney Foundation website, 2018. 13. Milliner DS. *Am J Nephrol.* 2005;25(2):154-160.